

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

EUROPEAN JOURNAL OF IMMUNOLOGY 5/1989

echa-
rcular
fane
ility in
l pre
sin A
mast
gene
vibro
acted
his
HIV
pen-
arar
the
vous
cted






EXHIBIT A

Shelley Schwarzbau[✉],
 Ahuva Nissim[○],
 Irit Alkalay[○],
 Mireille C. Ghori[○],
 Daniel G. Schindler[○],
 Yehudit Bergman[△] and
 Zelig Eshhar[○]

Department of Chemical Immunology,
 Weizmann Institute of Science[○],
 Rehovot and
 Department of Clinical Immunology,
 Hadassah Medical School[△],
 Jerusalem

Mapping of murine IgE epitopes involved in IgE-Fc_ε receptor interactions*

The generation of anti-IgE monoclonal antibodies has permitted the identification of various serological epitopes on the IgE molecule. The relationship of the sites on IgE recognized by such antibodies to the Fc epsilon receptor (Fc_εR) interaction site has been determined using cross-inhibition studies. However, interpretation of this type of experiment is limited by problems of steric hindrance. Thus, to accomplish precise mapping on the IgE molecule of the Fc_εR interaction site and the binding sites of various anti-IgE mAb, we employed site-directed mutagenesis of the IgE heavy chain gene.

To this end we have constructed and expressed a recombinant murine constant heavy chain (C_ε) gene bearing a (4-hydroxy-3-nitrophenyl)acetic acid (NP)-binding V_H region. Several site-specific mutants in the C_ε3 and C_ε4 domains of this recombinant C_ε gene were prepared and expressed by transfection into the light chain-producing J558L myeloma cell line. The resulting IgE antibodies were tested for binding to mast cells and to various anti-IgE mAb.

The mutants produced include a proline to histidine point mutant at amino acid residue 404 in the C_ε3 domain, a mutant with a truncated C_ε4 domain, a mutant with a 45 amino acid deletion in the carboxy end of C_ε3, and a chimeric human C_ε in which the human C_ε3 was replaced by the homologous mouse C_ε3 domain. These mutants have permitted the localization, to the C_ε3 domain, of the epitopes recognized by the 84.1C and 95.3 anti-IgE mAb. The 84.1C mAb recognizes a site on IgE which is identical or very close to the Fc_εR binding site, and 95.3 recognizes a site on IgE which is related, but not identical to the Fc_εR binding site. The antigenic determinant recognized by the 51.3 mAb, which is inefficient at blocking the IgE-Fc_εR interaction, has been mapped to the C_ε4 domain. When tested for binding to the Fc_εR on RBL-2H3 cells, the point mutant bound to the Fc_εR with twofold reduced affinity, while the C_ε3 deletion mutant and the mutant truncated in C_ε4 lost all receptor binding activity. These data suggest that the Fc_εR binding site can be assigned to the third C region domain, and that the fourth domain, while not directly involved in Fc_εR binding, may play a role in the formation of the H₂L₂ tetrameric IgE molecule, and in stabilizing the conformation of IgE required for Fc_εR binding.

1 Introduction

IgE antibodies are the class of antibody responsible for mediating the allergic response. Mast cells bear an Fc_ε receptor (Fc_εR) which is able to bind to IgE with high affinity. Antigen binding causes the cross-linking of the mast cell-bound IgE, triggering a series of responses in the mast cell leading to degranulation and the release of histamine and other mediators of allergy.

[I 7408]

* This research was supported by a grant from the National Council and Development, Israel, and the European Economic Community; Z.E. is an incumbent of the Marshall and Renette Ezralow Chair in Chemical and Cellular Immunology.

* Fellow of the Jane Coffin Childs Memorial Fund for Medical Research.

Correspondence: Zelig Eshhar, Department of Chemical Immunology, Weizmann Institute of Science, Rehovot 76100, Israel

Abbreviations: Fc_εR: Receptor for the constant region of IgE
 mAb: Monoclonal antibody(ies) NP: (4-Hydroxy-3-iodo-5-nitrophenyl)acetic acid NP: (4-Hydroxy-3-nitrophenyl)acetic acid
 ELISA: Enzyme-linked immunosorbent assay FgG: Fowl γ-globulin
 PBS: Phosphate-buffered saline

Much work has been done in an attempt to identify the site on the IgE molecule which interacts with the Fc_εR. Most of the data available to date implicate the third constant (C) region domain, and the interface between the second and the third domain of IgE, in receptor binding. Experiments by Pérez-Montfort and Metzger [1] have shown that a region in the cleft between the second and third C region domains is protected from proteolysis when bound to the Fc_εR. Burt et al. [2, 3] made synthetic peptides corresponding to regions of the rat IgE molecule thought to be accessible to the external environment. One of these peptides, P129, corresponding to residues 414–428 in the third C region domain (C_ε3), (residues 401–415 using the numbering system of Liu [4], the convention adopted in this report) inhibits the binding of IgE to the Fc_εR on a rat mast cell line (RBL-2H3). The binding of P129 is about 1000-fold less efficient on a molar basis than the binding of intact IgE.

In an analogous set of experiments, Robertson and Liu [5] have prepared antisera against peptides corresponding to regions of the mouse second, third, and fourth C region domains. Most of these antisera reacted against native IgE and showed varying reactivities against mast cell-bound IgE. One anti-peptide antibody, made against a peptide from C_ε3, "E-peptide-3", bound well to free IgE, but not to cell-bound IgE. This peptide may therefore represent a site sterically close to the epitope on IgE recognized by the Fc_εR. Interestingly, anti-

ethanol precipitated and used to transform competent bacteria (MC1061 strain).

2.3 Transfections and cell culture

RBL-2H3 cells [14], obtained from Drs. D. and T. Jovin, were grown in minimum Eagle's medium with the addition of 10% fetal calf serum (BioLab Ltd., Jerusalem, Israel), L-glutamine (Biological Industries, Kibbutz Beth Haemek, Israel), combined antibiotics (BioLab Ltd.) and fungizone (amphotericin, Biological Industries). J558L myeloma cells [15] (a gift of Dr. V. T. Oi), were grown in RPMI medium with 10% fetal calf serum (FCS), L-glutamine, and combined antibiotics (referred to as J558L medium).

Transfections were carried out by electroporation [16] or protoplast fusion [17]. For electroporation, 20 µg of purified plasmid was mixed with 2×10^7 J558L cells in phosphate-buffered saline (PBS) and subjected to a 1.8 kV discharge with an Isco (Nebraska, CO) power supply model 494. Cells were put into regular J558L medium for 24–48 h recovery and then transferred to selective medium.

Protoplast fusion was done as described [17]. Protoplasts prepared from 25-ml bacterial culture, containing the plasmid of interest were fused with polyethylene glycol (PEG) 1550 (Serva, Heidelberg, FRG) to 2×10^7 J558L cells. The cells were plated in 24-well plates at 10^5 /ml and cultured in J558L medium in the presence of antibiotics (100 µg/ml kanamycin or gentamycin (Sigma, St. Louis, MO)), to kill any remaining viable bacteria. Selective medium was added at 24–48 h and clones were visible in 2–3 weeks. Selective medium contained regular J558L medium with the addition of 1 µg/ml mycophenolic acid, 250 µg/ml xanthine and 15 µg/ml hypoxanthine (Sigma).

2.4 Immunochemical reagents

Rabbit anti-mouse IgE was made in our laboratory by immunizing rabbits with several mouse mAb of the IgE class. Antisera were preadsorbed on normal mouse serum-coupled Sepharose and then further purified on an IgE-Sepharose affinity column. These antibodies recognize the C_ε chain and also react with λ1 light (L) chains which are present on the SPE-7 monoclonal IgE used for immunization. Biotinylation was done by the method of Kendall et al. [18]. NIP-protein conjugates were made by the method of Brownstone et al. [19] and coupled to Sepharose 4B following cyanogen bromide activation.

2.5 IgE purification

Lysates were produced from non-secreting mutants (AFE), by suspending cells in PBS, 1 mM EDTA, 1 mM phenylmethyl sulfonyl fluoride (Sigma), 5 mM iodoacetamide and 0.2 U/ml aprotinin (Sigma). Cell suspensions were lysed either by freeze-thawing, sonication or the addition of 1% Nonidet-P40 (NP40).

Anti-NIP [(4-hydroxy-3-iodo-5-nitrophenyl)acetate] IgE antibodies were purified on NIP-ovalbumin (OVA)-Sepharose columns. J558L transfectoma supernatant (200–500 ml) or

10 ml of cell lysate from 5×10^8 cells, were passed over NIP-OVA-Sepharose columns. The columns were then washed with PBS containing 0.5% NP40 and then extensively washed with PBS. Elution was carried out with 5 M MgCl₂. Eluates were dialyzed extensively against PBS, aliquoted and frozen.

2.6 Immunoassays for IgE anti-NIP antibodies

IgE anti-NIP antibodies were detected by one of several variations of the enzyme-linked immunosorbent assay (ELISA) or radioimmunoassays (RIA) described here. For the RIA, 96-well polyvinylchloride V-shaped wells (Cooke Labs., Alexandria, VA) were coated with 50 µl antigen solution (10 µg/ml NIP₂DNP₂OVA), or anti-IgE, overnight at 4°C. Plates were washed with PBS containing 0.1% bovine serum albumin (BSA), and blocked with PBS containing 1% BSA to reduce nonspecific binding. IgE preparations (usually 50 µl transfectoma supernatant) were then added for 2 h at room temperature, followed by ¹²⁵I-labeled anti-IgE. Wells were washed, cut apart and counted in a gamma counter.

For ELISA assays, flat-bottom ELISA plates (Nunc, Roskilde, Denmark) were coated with either 10 µg/ml NIP-fowl γ-globulin (FgG), or a 1:10 dilution of serum-free supernatant from hybridoma cells secreting anti-mouse IgE mAb [7, 8], diluted in 0.06 M sodium carbonate buffer, pH 9.5. After overnight incubation at 4°C, the plates were washed in PBS containing 0.05% Tween, and dilutions of IgE-containing transfectoma supernatants, or purified antibody were added, and the plates were incubated for 2 h at 37°C. Plates were then washed and a 1:500 dilution of biotinylated rabbit anti-mouse IgE was added for 2 h. After washing, plates were treated for 30 min with avidin-coupled peroxidase (Bio-Makor, Rehovot, Israel), washed and peroxidase substrate was added. Alternatively, for the anti-NIP ELISA, after incubation with the IgE samples, plates were treated with a 1:1000 dilution of peroxidase-conjugated anti-mouse Fab (Bio-Makor). Peroxidase substrate was added after washing. The substrate used was 2,2'-azino-bis(3-ethylbenzthiazoline 6-sulfonic acid) (Sigma), dissolved at 1 mg/ml in 28 mM citric acid, 44 mM Na₂HPO₄, 0.3% H₂O₂. Assays were read in an automatic ELISA reader (Titertek Multiskan; Flow Labs., Rockville, MD) at 620 nm.

2.7 Binding of iodinated IgE or RBL-2H3 cells

Purified IgE derived from transfectomas was iodinated using the chloramine-T method. Protein (40–100 µg) in 100 µl PBS was reacted with 0.5–1.0 mCi (= 18.5–37 MBq) of ¹²⁵I and 10 µl of chloramine T (2 mg/ml) for 2 min. The reaction was stopped with 20 µl of sodium disulfide (2 mg/ml) and the mixture chromatographed on a 7-ml Sephadex G-25 column in the presence of 1% BSA, KI, and hemoglobin solution to color the protein peak. The first radioactive peak was collected, and incorporation of iodine checked by trichloroacetic acid precipitation.

For binding studies, 10⁶ RBL-2H3 cells/tube were mixed with dilutions of radiolabeled IgE, in 250 µl Tyrode's buffer [20] with 1 mM EDTA. For competition assays, cold IgE competitor was added and the mixture incubated for 2 h before the addition of the radiolabeled IgE. After an additional 2-h incubation with shaking at 4°C, 50-µl aliquots were removed in

triplicate and layered over horse serum in microtubes. The tubes were centrifuged in a microfuge for 30 s, supernatant was aspirated and pellets were counted in a gamma counter. An additional 50- μ l aliquot from each sample was counted to determine total counts added. Direct binding data were analyzed by Scatchard analysis and linear least square regression.

2.8 Degranulation assay

RBL-2H3 cells (10^6) were incubated with each IgE sample for 30 min at 37°C. Cells were washed with Tyrode's buffer and resuspended in 1 ml Tyrode's buffer. Aliquots (100 μ l) were placed in microtiter wells to which were added either 250 ng/ml NIP-FgG to measure specific release or 15 μ l 10% Triton to measure maximum release, or nothing to measure spontaneous release. The plates were incubated for 30 min at 37°C and then 15 μ l of supernatant was removed from each well to measure β -hexosaminidase release, an indicator of degranulation. The supernatants removed were mixed in wells of an ELISA plate with 40 μ l substrate solution [1.3 mg/ml paranitrophenyl-N-acetyl- β -D-glucosamine (Sigma) in 0.04 M phosphate/citrate buffer, pH 4.5] and incubated 1–2 h at 37°C. To stop the reaction 0.2 M glycine, pH 10.7, was added and wells were read in an ELISA reader at 405 nm. Percent specific release was calculated as:

$$\frac{(\text{specific release} - \text{spontaneous release})}{(\text{maximum release})} \times 100,$$

where spontaneous release is release in the absence of antibody, and maximum release is the β -hexosaminidase activity obtained after 1% Triton lysis of cells.

2.9 Adsorption assay for binding to the Fc γ R

As an alternative means to determine the extent of binding of the mutant IgE to the Fc γ R on mast cells, we measured the ability of RBL-2H3 cells to remove anti-NIP activity from IgE-containing preparations. Dilutions (100 μ l) of purified mutant and wild-type IgE were incubated in triplicate in microtiter wells with either 100 μ l PBS or 100 μ l PBS containing 10^6 RBL-2H3 cells. The plates were incubated for 3–4 h at room temperature, with periodic shaking. Supernatant (100 μ l) was then removed from each well, and assayed for anti-NIP activity by ELISA, on NIP-FgG-coated ELISA plates. The ratio of optical density obtained from the ELISA from IgE preparations incubated with RBL-2H3 cells vs. those incubated without RBL-2H3 cells gives a measure of how much IgE was removed by these cells, and thereby indicates to what extent the IgE was able to bind to these cells.

2.10 Western blots and immunodetection

IgE transfectoma lysates, cell supernatants, or purified IgE preparations were run on sodium dodecyl sulfate (SDS) polyacrylamide gels [21]. After electrophoresis proteins were transferred to nitrocellulose in 15.6 mM Tris, 120 mM glycine, pH 8.3 for 4 h at 200 mA. The nitrocellulose filters were stained with Ponceau red (Sigma) to visualize protein standards, prehybridized in PBS containing 10% skim milk for 16 h at 4°C and then hybridized in the above buffer with 125 I-labeled rabbit anti-mouse IgE. Filters were washed in PBS and PBS-Tween and autoradiographed.

3 Results

3.1 Construction of the PSV2-V μ -C μ plasmid

In order to generate IgE molecules with various mutations in the Fc portion of their C region, we constructed a vector which permitted the expression, in a mammalian cell, of a complete IgE molecule of known antigenic specificity. This was accomplished by joining the gene segment from a genomic clone encoding the mouse C μ region with a V μ gene fragment encoding a V region with anti-NP specificity [9]. The V region encoded by this gene is heteroclitidic and binds the antigen NIP with higher affinity than NP. All assays measuring hapten binding were therefore performed using NIP-conjugated rather than NP proteins. The PSV2-V μ -C μ plasmid which we constructed (see below) contains the entire mouse C μ coding sequence and an anti-NP variable region (Fig. 1C). Upon transfection into the L chain-producing J558L myeloma cell line, the ϵ chain encoded by the recombinant gene combines with the endogenous J558L L chain to produce an IgE antibody with anti-NP specificity. Similar constructs have been used previously by Neuberger et al. [22] to obtain chimeric IgE molecules bearing a human ϵ gene and a NP-binding mouse V region.

The PSV2-V μ -C μ plasmid was constructed from the PSV2-V μ 6 vector [9] and Fig. 1A). This expression vector contains a rearranged V μ gene of anti-NP specificity [9]. The 6.25-kb fragment obtained after BamHI digestion was ligated to the 4.6-kb BamHI fragment of the PBR-C μ plasmid (Fig. 1B), containing the full mouse C μ region [11]. The vector so obtained (denoted PSV2-C μ -E) contained the full C μ and V μ segment but was lacking the enhancer element for Ig transcription. In order to increase the expected levels of Ig expression with this vector, the 1-kb fragment containing Ig enhancer activity, obtained from XbaI digestion of the pA-6 plasmid, was inserted into the unique EcoRI site of the PSV2-C μ -E vector by blunt-end ligation of filled-in ends. The vector thus obtained was called PSV2-V μ -C μ and its structure is shown in Fig. 1C.

The PSV2-V μ -C μ plasmid was introduced into J558L L chain-producing myeloma cells using the technique of protoplast




Figure 2. Immunoblot analysis of mutant and wild-type IgE antibodies. Purified IgE preparations (25–100 ng/lane) were run under reducing (A) or nonreducing (B) conditions through 10% or 5%–10% gradient acrylamide gels, respectively, electroblotted to nitrocellulose and hybridized with iodinated rabbit anti-mouse IgE antibodies. Note that in addition to the ϵ chain, the anti-IgE antibodies cross-react with the λ L chain.

Table 1. Properties of ITC5 recombinant IgE

Property	ITC5	SPE-7
Chain apparent molecular weight	25 kDa	25 kDa
K _d for Fc ϵ R binding	2.5 $\times 10^{-8}$ M	2.5 $\times 10^{-8}$ M
Fc ϵ R binding inhibition	50% at 100 ng/ml	50% at 100 ng/ml
Degranulation of RBL-2H3 cells	50%	50%

- Mouse monoclonal IgE [23].
- Determined by Scatchard analysis of binding of radioiodinated IgE to RBL-2H3 cells.
- Concentration of IgE giving 50% inhibition of the binding of radioiodinated SPE-7 to RBL-2H3 cells.
- % RBL-2H3 cell degranulation induced by IgE at 100 ng/ml, triggered by NIP-DNP-BSA.

fusion. Supernatants from clones surviving growth in selective medium containing mycophenolic acid were screened for IgE anti-NIP activity using a RIA (not shown) and the ITC5 clone was chosen for all further characterization. This clone secretes approximately 2 μ g IgE/ml. Table 1 summarizes properties of the ITC5 IgE as compared to SPE-7 hybridoma-derived monoclonal IgE [23]. Both IgE have similar molecular weights (Fig. 2), bind with similar affinities to the Fc ϵ R on mast cells and both are able to induce RBL-2H3 degranulation in the presence of cross-linking antigen.

3.2 Generation of vectors encoding mutant IgE antibodies

The mutated IgE antibodies that we describe in this study are depicted schematically in Fig. 3 and Table 2. They include AFE, truncated in C₄, DSE, with a 45-amino acid deletion in the C₃ domain, and PHS, with a proline to histidine point mutation in C₃. The AFE mutant was made by creating a frameshift mutation at the Aval site in the C₄ domain (Fig. 1B, C). This frameshift caused a stop codon to be reached at amino acid 472 (position 34 of the C₄ domain). The region between the mutation site and stop codon was expected to have altered sequence due to the frameshift introduced (Fig. 3).

To generate this mutation, the PSV2-V_H-C₂ vector was partially digested with Aval and the 11.85-kb linearized fragment was blunt ended with the Klenow fragment of DNA polymerase I. This was self ligated and used to transform bacteria. Restriction mapping and DNA sequencing were used to confirm that the correct Aval site was mutated (not shown). The vector containing the AFE mutation was used to transfect J558L cells. Clones surviving mycophenolic acid selection were tested by an ELISA assay for secretion of anti-NIP antibody. Of nine clones tested from two separate transfection experiments, none secreted detectable amounts of IgE. (The sensitivity of the ELISA is approximately 5 ng/ml). It is possible that the cells can produce IgE, but that due to the gross structural alteration of IgE, they are unable to secrete IgE. We therefore produced cell lysates, using sonication, freeze-thawing or NP40 lysis. The lysates were tested for IgE anti-NIP activity by ELISA and were found to contain between <5 and 110 ng/10⁶ cells. Immunoblot analysis (Fig. 2) under reducing conditions of affinity-purified AFE mutant IgE from such lysates revealed H chains of reduced size as expected for this mutant. Electrophoresis under nonreducing conditions reveals

that almost all of the antibody is present as HL, instead of the full H₂L₂ tetramer (Fig. 2).

Another deletion mutant (DSE) was constructed with a deletion of 45 amino acids in C₃ and the beginning of C₄ (amino acid residues 397-441). The deletion encompasses the sequence spanned by the P129 peptide of Burt and Stanworth [2] and also includes a cysteine which participates in the C₃ intradomain disulfide bridge [4, 24]. This mutant was made by digestion of the PSV2-V_H-C₂ plasmid with Spe I and Eco 47III (each of which cut the plasmid at a single site), blunting the ends with mungbean nuclease, religating the large linear fragment and transforming bacteria with the product of this reaction. The structure of this plasmid was checked by restriction mapping and by hybridization to an oligomer which was complementary to the expected sequence of the junction site after ligation. Transfected clones were found to secrete antibody, but in very small amounts, 5-40 ng/ml.

A proline to histidine point mutation (PHS) was created at amino acid residue 404, using oligonucleotide-directed, site-

	C ₂	S-S (L)
HOUSE C ₂ (ITC5)	SH NPGLYFLKPS	KGTASHTLGG
PHS (P to H point mutation)	SH NPGLYFLKPS	KGTASHTLGG
AFE (C ₄ frameshift)	SH NPGLYFLKPS	KGTASHTLGG
DSE (deletion in C ₃)	SH NPGLYFLKPS	KGTASHTLGG
	CHO	
HOUSE C ₂	TUTUYDELM	STVYVFALE
PHS	TUTUYDELM	STVYVFALE
AFE	TUTUYDELM	STVYVFALE
DSE	TUTUYDELM	STVYVFALE
	CHO	S-S (H)
HOUSE C ₂	HESTTLVRF	VMTPELEL
PHS	HESTTLVRF	VMTPELEL
AFE	HESTTLVRF	VMTPELEL
DSE	HESTTLVRF	VMTPELEL
	CHO	
HOUSE C ₂	SULMDDREIT	DTLAQTVLM
PHS	SULMDDREIT	DTLAQTVLM
AFE	SULMDDREIT	DTLAQTVLM
DSE	SULMDDREIT	DTLAQTVLM
	(H)-S-S	C ₃
HOUSE C ₂	SQCVOYLANT	NACFONEFEG
PHS	SQCVOYLANT	NACFONEFEG
AFE	SQCVOYLANT	NACFONEFEG
DSE	SQCVOYLANT	NACFONEFEG
	CHO	CHO
HOUSE C ₂	RENVNVTUNG	ENKTSVASAQ
PHS	RENVNVTUNG	ENKTSVASAQ
AFE	RENVNVTUNG	ENKTSVASAQ
DSE	RENVNVTUNG	ENKTSVASAQ
	C ₄	
HOUSE C ₂	QCIVNDPDPF	KDIVSITKT
PHS	QCIVNDPDPF	KDIVSITKT
AFE	QCIVNDPDPF	KDIVSITKT
DSE	QCIVNDPDPF	KDIVSITKT
	CHO	CHO
HOUSE C ₂	YFFPDIEVQ	VLCOCKLISN
PHS	YFFPDIEVQ	VLCOCKLISN
AFE	YFFPDIEVQ	VLCOCKLISN
DSE	YFFPDIEVQ	VLCOCKLISN
	CHO	CHO
HOUSE C ₂	VTQREKPTQR	VIRHALQKPR
PHS	VTQREKPTQR	VIRHALQKPR
AFE	VTQREKPTQR	VIRHALQKPR
DSE	VTQREKPTQR	VIRHALQKPR

Figure 3. Amino acid sequence of the mouse C₂ domains and of the IgE mutants prepared in this study. Dashes (—) indicate wild-type sequence; (x) indicates deleted amino acids; (CHO) indicates glycosylation sites and (S-S) indicates intrachain disulfide bonds. S-S (H)- and S-S (L)- represent the interchain disulfide bonds with H and L chains. The sequence shown is based on references [4] and [23] and the one-letter amino acid code is given in [4].

Table 2. Summary of mutants produced to this study

a) All the recombinant IgE molecules listed above contain a NIP-specific mouse V_H region with a C_ϵ region and express the endogenous J558L mouse λ 1 L chain.

specific mutagenesis [13]. This mutation is located in $C_{\epsilon 3}$, in the region spanned by the P129 oligomer of Burt and Stanworth [2]. Aliquots of the PBR- C_ϵ plasmid were digested with SalI and EcoRV, and the resulting digests were mixed, melted, and reannealed in the presence of a molar excess of kinased SAS3 oligomer (sequence GGC, AAC, TAC, ATG, TAG, GAT, GGA). This oligomer is complementary to the sequence encoding amino acids 401-407, with a C to A mutation which alters proline 404 to histidine. The oligomer also contains a silent mutation to increase the ease of detecting mutant vs. wild type. The positions of the two mutations are underlined. Mutant colonies were identified by hybridization with radiolabeled oligomer. The 4.6-kb insert corresponding to C_ϵ was removed by digestion with BamHI and ligated into the PSV2- V_H - C_ϵ plasmid from which the wild-type C_ϵ sequences were removed. This plasmid (PSV- V_H - C_ϵ -PHS) was used to transfect J558L cells by electroporation. Supernatants from clones surviving mycophenolic acid selection were tested for IgE anti-NIP antibodies. The PHS clone selected secretes approximately 2 μ g/ml of antibody. RNA was isolated from this clone and tested by Northern blot hybridization with the mutant oligomer for the presence of the mutated sequence. The oligomer hybridizes to PHS but not to ITC5 (wild-type) RNA (not shown). SDS gel electrophoresis of the IgE produced by the PHS mutant reveals H and L chains of the expected size, present as H_2L_2 in nonreduced gels (Fig. 2).

The preparation and expression of vectors containing the human C_ϵ sequences, and human-mouse chimeras will be described in a separate manuscript (A. Nissim and Z. Eshhar, in preparation).

3.3 Binding of the mutant IgE to the $Fc_\epsilon R$ on RBL-2H3 cells

In order to identify the sites on the IgE molecule involved in the interaction with the $Fc_\epsilon R$, the IgE mutants were tested for their ability to bind to RBL-2H3 cells, a rat mast cell line bearing high-affinity $Fc_\epsilon R$ for rodent IgE [14]. A number of different assays were used to measure binding. First, affinity-purified preparations of unmutated ITC5 and the IgE mutants were tested for their ability to inhibit the binding of iodinated, purified monoclonal IgE (SPE-7) [24] to RBL-2H3 cells. The inhibition curves so obtained are shown in Fig. 4A. It may be seen that the PHS point mutant is reduced approximately twofold in its ability to bind to the $Fc_\epsilon R$, while the DSE mutant has lost all ability to compete with radioiodinated IgE for the binding to RBL-2H3 cells. The AFE mutant, which is not secreted, and was therefore purified from cell lysates, was not

available in sufficient concentrations for the inhibition assay. It was therefore tested in the other binding assays described below. Since the differences in the inhibition of binding by the ITC5 wild-type IgE and the PHS mutant were small, these IgE preparations were also tested for direct binding to RBL-2H3 cells. Affinity-purified ITC5 and PHS were radioiodinated and the binding of these antibodies to RBL-2H3 cells was tested. Scatchard analysis (Fig. 4B) reveals that the PHS mutant has a 2.5-fold lower affinity for the $Fc_\epsilon R$ than ITC5 wild-type IgE.

These results were further confirmed in two other assays. In an assay measuring the degranulation of RBL-2H3 cells, the ITC5 (wild type) and PHS (mutant) IgE antibodies, but not the DSE or AFE mutant IgE were able to induce degranulation of RBL-2H3 cells, when cross-linked on the cell surface by polyvalent antigen (NIP-FgG) (Table 3).




Figure 4. $Fc_\epsilon R$ binding of mutant IgE antibodies. (A) Inhibition of binding of iodinated IgE to RBL-2H3 cells by mutant and wild-type recombinant IgE. RBL cells were preincubated with varying concentrations of ITC5 (wild type), PHS (point mutant) or DSE (deletion mutant) IgE. 125 I-labeled monoclonal IgE was then added and after a 2-h incubation, cell-bound and free IgE were separated and radioactivity was counted. All determinations were done in triplicate. (B) Scatchard analysis of binding of ITC5 (unmutated) and PHS (point mutant) IgE. RBL-2H3 cells were incubated with several concentrations of iodinated ITC5 and PHS IgE. Unbound counts were removed by centrifugation through serum, and the cell pellet was counted. Lines represent the least squares fit, derived by linear regression analysis.

Table 3. Degranulation of RBL-2H3 cells mediated by IgE mutants

IgE mutant	NIP-FcG	β -Hexosaminidase release
ITC5	High	High
PHS	Low	Low
AFE	Low	Low
DSE	High	High
NO IgE	None	None

- a) RBL-2H3 cells were preincubated with affinity-purified IgE preparations at 0.2 μ g/ml for 30 min and washed before the addition of NIP-FcG (250 ng/well). β -Hexosaminidase release was determined as described in Sect. 2.8. Each determination was done in triplicate. Standard deviations were < 15% of ELISA readings.

Finally, mutant IgE antibodies were tested for their ability to be adsorbed by RBL-2H3 cells. Dilutions of the affinity-purified antibodies were incubated with RBL-2H3 cells to allow binding. Supernatants containing unbound IgE were tested by ELISA for anti-NIP activity. Results were plotted as the ratio between the ELISA readings obtained for adsorbed supernatants, divided by unadsorbed supernatants, vs. the concentration of the IgE preparation used (represented by ELISA reading). As can be seen in Fig. 5, the ratio of A_{420} "adsorbed"/ A_{420} "unadsorbed" was lowest for ITC5, the wild-type IgE (best adsorbed), and higher for the PHS mutant, though some anti-NIP activity was still removed. For the AFE and DSE mutants, the ratio was close to one, indicating that the RBL-2H3 cells did not remove any of the IgE anti-NIP antibodies. Thus, we may conclude that binding of mutant IgE molecules to the $Fc\gamma R$ on rat mast cells follows the order: ITC5 > PHS > AFE = DSE.

3.4 Recognition of mutant IgE antibodies by anti-IgE mAb

Purified mutant and wild-type IgE antibodies were next tested for their ability to bind to anti-mouse IgE mAb which have been previously described [7, 8]. ELISA assays were performed in which wells were coated with either antigen (NIP-FcG), or serum-free preparations of the anti-IgE mAb. This was followed by dilutions of purified wild-type and mutant IgE preparations. Degree of binding was determined using biotinylated rabbit anti-mouse IgE (H+L), and peroxidase-labeled avidin. Table 4 shows the results of one representative assay.

Table 4. Binding of IgE mutants to anti-IgE mAb

IgE mutant	Anti-mouse IgE mAb
ITC5	High
PHS	Low
AFE	Low
DSE	High
NO IgE	None

lated rabbit anti-mouse IgE (H+L), and peroxidase-labeled avidin. Table 4 shows the results of one representative assay.

It may be seen that the PHS point mutant has lost the ability to be recognized by the 95.3 anti-IgE mAb, while it is still recognized by the 51.3 and 84.1C mAb (Table 4) and can still bind to mast cells (though with reduced affinity) (Fig. 4). The DSE deletion mutant has lost both recognition by all three mAb and the ability to bind to the $Fc\gamma R$ on mast cells. The AFE mutant which has a frameshift causing premature termination in the C₄ domain has reduced binding to the 51.3 anti-IgE mAb, and has lost the ability to bind to the $Fc\gamma R$ on RBL-2H3 cells. A chimeric human IgE (H₁MC₃) containing mouse C₃ (manuscript in preparation), binds the 84.1C and 95.3 mAb, but not the 51.3 mAb. "TAN" human IgE (manuscript in preparation), with the J558L L chain and mouse anti-NIP V_H region, is not recognized by any of the mAb. These data permit the assignment of the site recognized by 84.1C and 95.3 to the C₃ domain, and that recognized by 51.3 to the C₄ domain.

4 Discussion

In order to learn more about the $Fc\gamma R$ binding site on the IgE molecule, we have developed an *in vitro* expression system enabling the production of genetically engineered mouse IgE with anti-NIP activity. The antibody so produced, represented by clone ITC5, has all the properties of authentic mouse IgE and binds with high affinity to the $Fc\gamma R$ on mast cells (Table 1). The plasmid encoding the ITC5 antibody was then subjected to a variety of mutagenesis procedures in order to produce IgE molecules with alterations in their Fc region. These mutants were used in binding assays to determine their ability to bind to the $Fc\gamma R$ on mast cells and to bind to a series of anti-IgE mAb previously generated in our laboratory [7, 8].

Four assays were employed to measure the binding of mutant or wild-type IgE to the $Fc\gamma R$ on mast cells: (a) the inhibition of binding of radiolabeled IgE to RBL-2H3 cells, (b) the direct binding of iodinated IgE to RBL-2H3 cells, (c) IgE-mediated degranulation of RBL cells and (d) the adsorption, by RBL-2H3 cells, of anti-NIP activity from IgE-containing solutions. These assays gave concordant results. Compared to the ITC5 wild-type IgE, the PHS proline to histidine point mutant bound the $Fc\gamma R$ with approximately twofold reduced affinity (Figs. 4A, B and 5). PHS mutant IgE induced almost as much degranulation as did ITC5 wild-type IgE (Table 3). Since only

- Purified unmutated (ITC5) and mutated recombinant IgE were tested by ELISA for binding to various anti-mouse IgE mAb. ELISA plates were first coated with serum-free supernatant containing the various anti-IgE mAb, or with NIP-conjugated protein. IgE (4 ng) diluted in medium (DMEM and FCS) was then added to each well. After washing, the wells were treated with biotin-conjugated rabbit anti-mouse IgE, followed by an avidin-peroxidase conjugate and peroxidase substrate. Absorbance was read in an automatic ELISA reader. Variation between duplicates was always < 10% of the values read, or < 0.01 absorbance units for negative wells.
- Anti-mouse IgE mAb.
- See Table 2.
- Mouse C₃ into human C₄1,2,3.
- Human IgE.




Figure 5. Adsorption of IgE by RBL-2H3 cells. Dilutions of mutant and wild-type IgE antibodies were incubated with RBL-2H3 cells for 4 h in microtiter wells. After this incubation, supernatants were removed and tested by ELISA for anti-NIP activity. Results were plotted as the adsorption index (the ratio between the ELISA readings obtained for adsorbed divided by unadsorbed supernatants) vs. the concentration of the IgE preparation tested. An adsorption index closer to unity indicates less adsorption of IgE by RBL-2H3 cells, and hence, less binding of the particular mutant.

a small fraction of $Fc_\gamma R$ on RBL-2H3 cells need to be occupied in order to induce degranulation, it is not surprising that sufficient occupancy could be obtained with either ITC5 wild type or PHS point mutant IgE.

The AFE frame shift mutant, lacking a functional $C_\gamma 4$ domain, and the DSE mutant, with a 45-amino acid deletion in $C_\gamma 3$, are unable to bind to the $Fc_\gamma R$ in the four assays used (Figs. 4 and 5 and Table 3). It is surprising, in light of the accumulated evidence [1-3, 5] suggesting that the binding site for the $Fc_\gamma R$ is present on $C_\gamma 3$, that the AFE mutant (lacking the $C_\gamma 4$ domain) is unable to bind to the $Fc_\gamma R$. We believe that the $C_\gamma 4$ domain, while not containing the actual receptor binding site(s), is necessary for the IgE molecule to attain the correct conformation needed for receptor binding. The $C_\gamma 4$ domain has been shown to be important for the dimerization of human IgE [6]. Interestingly, our AFE mutant mouse IgE which lacks most of the $C_\gamma 4$ domain, is present inside the cell almost entirely as HL (Fig. 2). This indicates that in mouse IgE, the fourth domain may also have a role in initiating, or stabilizing H chain dimerization.

The epitopes recognized by three anti-IgE mAb, 84.1C, 95.3 and 51.3, were assigned to domains on the ϵ chain based on their ability to recognize the various IgE mutants tested. For example, the 95.3 mAb was shown to recognize a determinant on the $C_\gamma 3$ domain, since it is no longer recognized by the PHS mutant, which has a single proline to histidine substitution in $C_\gamma 3$, and is still recognized by the AFE mutant which has 96% of its $C_\gamma 4$ domain altered or deleted. In addition, 95.3 is able to recognize the mouse-human chimeric IgE, which has only mouse $C_\gamma 3$, while it is unable to recognize human IgE. By analogous logic, the epitope recognized by 84.1C may be assigned to $C_\gamma 3$, and the epitope recognized by 51.3 is most likely on $C_\gamma 4$ (Table 4).

The purified DSE mutant (45-amino acid deletion in $C_\gamma 3$) seems to bind none of the anti-IgE mAb, including the 51.3 antibody which we have assigned to $C_\gamma 4$. However, when DSE transfectoma supernatant was tested in a similar assay, some

reaction was seen against the 51.3 and 84.1C mAb (not shown). It is possible that conformational changes in this mutant have reduced binding to these two anti-IgE mAb, and that these conformational changes in the DSE mutant are more pronounced when the mutant IgE is subjected to high-salt elution from the affinity column. Similarly, in a number of experiments, when fresh DSE supernatants were tested for binding to the $Fc_\gamma R$ on RBL cells using the adsorption assay, some binding to the $Fc_\gamma R$ was detected. Purified fractions of this mutant, eluted with 5 M $MgCl_2$, consistently failed to bind. Thus, the effects of the deletion on the binding to the 84.1C and 51.3 anti-IgE mAb, and on the binding to the $Fc_\gamma R$ are likely to be conformational, rather than due to the elimination of the binding site. We have previously shown [7, 8] that the 51.3, 95.3 and 84.1C anti-IgE mAb are sensitive to IgE conformation and are unable to bind to denatured or reduced IgE.

The anti-IgE antibodies have previously been analyzed in inhibition studies to determine the relationship of the epitopes they recognize to the binding site for IgE. The 51.3 antibody inhibits poorly the binding of IgE to its receptor and is able to bind to IgE receptor complexes [7]. Thus, it seems that this antibody recognizes an epitope which is not closely related to the receptor binding site. This is consistent with our conclusion that the epitope recognized by this antibody is on the $C_\gamma 4$ domain. The 95.3 anti-IgE mAb inhibits the binding of IgE to its $Fc_\gamma R$ but can still bind to cell-bound IgE [7]. Thus the epitope recognized by this antibody is related, but not identical to the receptor binding site. This is also apparent from the fact that the PHS point mutant has lost, entirely, the epitope recognized by 95.3, but is still able to bind to the $Fc_\gamma R$ on mast cells, albeit with reduced affinity. The reduced ability of the PHS mutant to bind to the $Fc_\gamma R$ on RBL cells may be due to a direct effect of the proline to histidine mutation in destabilizing the IgE- $Fc_\gamma R$ interaction, or could be due to conformational changes caused by this mutation which are propagated to distal sites on the $C_\gamma 3$ domain. While more mutants will be required in order to distinguish between these possibilities, it is clear that the 95.3 binding site can be totally abolished with only a moderate effect on $Fc_\gamma R$ binding.

The 84.1C anti-IgE mAb inhibits the binding of IgE to the $Fc_\gamma R$ and does not recognize cell-bound IgE [8]. Thus, it seems likely that this antibody recognizes an epitope very closely related to the binding site for the $Fc_\gamma R$. It will be interesting to see whether among further IgE point mutants which we are currently producing, one will be found which loses both the ability to bind to the $Fc_\gamma R$ and to the 84.1C anti-IgE mAb. A good candidate for such a mutation may be a site on the "E-peptide-3" produced by Liu et al. [5]. Antibodies against this $C_\gamma 3$ peptide behave like our 84.1C mAb in that they bind to free, but not $Fc_\gamma R$ -bound IgE, and thus seem to cross-inhibit the IgE- $Fc_\gamma R$ interaction.

By combining the above data, a few conclusions may be reached. We have mapped to the $C_\gamma 3$ domain the 84.1C anti-IgE mAb most closely related to the IgE- $Fc_\gamma R$ interaction site. This determinant is not included within the P129 peptide site described by Burt and Stanworth [2]. A single proline to histidine substitution in the region spanned by the P129 peptide leads to the loss of recognition by the 95.3 mAb, which recognizes an epitope that is related, but not identical to the $Fc_\gamma R$ binding site. However, such a mutant (PHS) retains full recognition by the 84.1C anti-IgE mAb, and exhibits only two-fold reduction in binding to the $Fc_\gamma R$. Thus, the $Fc_\gamma R$ binding

site is probably located in the C₃ domain, but not in the region encompassed by P129. The C₄ domain seems to be required for the formation of the H₂L₂ tetramer and, though not directly involved in Fc_εR binding, may play a role in stabilizing the conformation of IgE required for binding to the Fc_εR. The generation of more IgE deletion and point mutants, together with mouse/human exon shuffling experiments (manuscript in preparation) should allow the precise identification of sites on the IgE molecule required for dimerization, binding to mAb and binding to the high-affinity Fc_εR on mast cells.

The authors wish to thank Ms. S. Barak and Ms. T. Waks for technical assistance, and Drs. T. Honjo, D. Rice and V. Oi for the gifts of plasmids and cell lines.

Received January 16, 1989; in revised form March 13, 1989.

5 References

- 1 Pérez-Montfort, R. and Metzger, H., *Mol. Immunol.* 1982, 19: 1113.
- 2 Burt, D. S. and Stanworth, D. R., *Eur. J. Immunol.* 1987, 17: 437.
- 3 Burt, D. S., Hastings, G. Z., Healy, J. and Stanworth, D. R., *Mol. Immunol.* 1987, 24: 379.
- 4 Liu, F.-T., Albrandt, F., Sutcliffe, J. G. and Katz, D. H., *Proc. Natl. Acad. Sci. USA* 1982, 79: 7852.
- 5 Robertson, M. W. and Liu, F.-T., *Mol. Immunol.* 1988, 25: 103.
- 6 Helm, B., Marsh, P., Vercelli, D., Padlan, E., Gould, H. and Geha, R., *Nature* 1988, 331: 180.
- 7 Baniyash, M. and Eshhar, Z., *Eur. J. Immunol.* 1984, 14: 799.
- 8 Baniyash, M., Kerhy, M. and Eshhar, Z., *Mol. Immunol.* 1988, 25: 705.
- 9 Bothwell, A. L. M., Pastind, M., Reth, M., Imanishi-Kari, T., Rajewsky, K. and Baltimore, D., *Cell* 1981, 24: 625.
- 10 Mulligan, R. C. and Berg, P., *Proc. Natl. Acad. Sci. USA* 1981, 78: 2072.
- 11 Ishida, N., Ueda, S., Hayashida, H., Miyata, T. and Honjo, T., *EMBO J.* 1982, 1: 1117.
- 12 Maniatis, T., Fritsch, E. F. and Sambrook, J., *Molecular Cloning. A laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 1982.
- 13 Morinaga, Y., Franceschini, T., Inouye, S. and Inouye, M., *Biotechnology* 1984, 2: 636.
- 14 Barsumian, E. L., Isersky, C., Petrino, M. G. and Siraganian, R. P., *Eur. J. Immunol.* 1981, 11: 317.
- 15 Oi, V. T., Morrison, S. L., Herzenberg, L. A. and Berg, P., *Proc. Natl. Acad. Sci. USA* 1983, 80: 825.
- 16 Porter, H., Weir, L. and Leder, P., *Proc. Natl. Acad. Sci. USA* 1984, 81: 7161.
- 17 Sandri-Goldin, R. M., Goldin, A. L., Levine, M. and Gironzo, J. C., *Mol. Cell Biol.* 1981, 1: 743.
- 18 Kendall, C., Ionescu-Maniu, I. and Dreesman, G. R., *J. Immunol. Methods* 1983, 56: 329.
- 19 Brownstone, A., Mitchison, N. A. and Ritt-Rivers, R., *Immunology* 1966, 10: 465.
- 20 Baniyash, M., Eshhar, Z. and Rivnay, B., *J. Immunol.* 1986, 136: 588.
- 21 Laemmli, U. K., *Nature* 1971, 227: 680.
- 22 Neuberger, M. S., Williams, G. T., Mitchell, E. B., Jouhal, S. S., Flanagan, J. G. and Rabbits, T. H., *Nature* 1985, 314: 268.
- 23 Eshhar, Z., Ofarim, M. and Waks, T., *J. Immunol.* 1980, 124: 775.
- 24 Ishida, N., Ueda, S., Hayashida, H., Miyata, T. and Honjo, T., *EMBO J.* 1982, 1: 1117.